

UNCERTAINTY

CHAPTER 13

Chapter 13 1

Outline

- ◇ Uncertainty
- ◇ Probability
- ◇ Syntax and Semantics
- ◇ Inference
- ◇ Independence and Bayes' Rule

Chapter 13 2

Uncertainty

Let action A_t = leave for airport t minutes before flight
Will A_t get me there on time?

Problems:

- 1) partial observability (road state, other drivers' plans, etc.)
- 2) noisy sensors (KCBS traffic reports)
- 3) uncertainty in action outcomes (flat tire, etc.)
- 4) immense complexity of modelling and predicting traffic

Hence a purely logical approach either

- 1) risks falsehood: " A_{25} will get me there on time"
- or 2) leads to conclusions that are too weak for decision making:
" A_{25} will get me there on time if there's no accident on the bridge
and it doesn't rain and my tires remain intact etc etc."

(A_{1440} might reasonably be said to get me there on time
but I'd have to stay overnight in the airport ...)

Chapter 13 3

Methods for handling uncertainty

Default or nonmonotonic logic:

Assume my car does not have a flat tire

Assume A_{25} works unless contradicted by evidence

Issues: What assumptions are reasonable? How to handle contradiction?

Rules with fudge factors:

$A_{25} \mapsto_{0.3} \text{AtAirportOnTime}$

$\text{Sprinkler} \mapsto_{0.99} \text{WetGrass}$

$\text{WetGrass} \mapsto_{0.7} \text{Rain}$

Issues: Problems with combination, e.g., *Sprinkler causes Rain??*

Probability

Given the available evidence,

A_{25} will get me there on time with probability 0.04

Mahaviracarya (9th C.), Cardano (1565) theory of gambling

(Fuzzy logic handles **degree of truth** NOT uncertainty e.g.,

WetGrass is true to degree 0.2)

Chapter 13 4

Probability

Probabilistic assertions **summarize** effects of

laziness: failure to enumerate exceptions, qualifications, etc.

ignorance: lack of relevant facts, initial conditions, etc.

Subjective or **Bayesian** probability:

Probabilities relate propositions to one's own state of knowledge

e.g., $P(A_{25} | \text{no reported accidents}) = 0.06$

These are **not** claims of a "probabilistic tendency" in the current situation
(but might be learned from past experience of similar situations)

Probabilities of propositions change with new evidence:

e.g., $P(A_{25} | \text{no reported accidents, 5 a.m.}) = 0.15$

(Analogous to logical entailment status $KB \models \alpha$, not truth.)

Chapter 13 5

Making decisions under uncertainty

Suppose I believe the following:

$P(A_{25} \text{ gets me there on time} | \dots) = 0.04$

$P(A_{90} \text{ gets me there on time} | \dots) = 0.70$

$P(A_{120} \text{ gets me there on time} | \dots) = 0.95$

$P(A_{1440} \text{ gets me there on time} | \dots) = 0.9999$

Which action to choose?

Depends on my **preferences** for missing flight vs. airport cuisine, etc.

Utility theory is used to represent and infer preferences

Decision theory = utility theory + probability theory

Chapter 13 6

Probability basics

Begin with a set Ω —the **sample space**

e.g., 6 possible rolls of a die.

$\omega \in \Omega$ is a **sample point/possible world/atomic event**

A **probability space** or **probability model** is a sample space with an assignment $P(\omega)$ for every $\omega \in \Omega$ s.t.

$$0 \leq P(\omega) \leq 1$$

$$\sum_{\omega} P(\omega) = 1$$

e.g., $P(1) = P(2) = P(3) = P(4) = P(5) = P(6) = 1/6$.

An **event** A is any subset of Ω

$$P(A) = \sum_{\{\omega \in A\}} P(\omega)$$

E.g., $P(\text{die roll} < 4) = P(1) + P(2) + P(3) = 1/6 + 1/6 + 1/6 = 1/2$

Chapter 13 7

Random variables

A **random variable** is a function from sample points to some range, e.g., the reals or Booleans

e.g., $Odd(1) = true$.

P induces a **probability distribution** for any r.v. X :

$$P(X = x_i) = \sum_{\{\omega: X(\omega) = x_i\}} P(\omega)$$

e.g., $P(Odd = true) = P(1) + P(3) + P(5) = 1/6 + 1/6 + 1/6 = 1/2$

Chapter 13 8

Propositions

Think of a proposition as the event (set of sample points) where the proposition is true

Given Boolean random variables A and B :

event a = set of sample points where $A(\omega) = true$

event $\neg a$ = set of sample points where $A(\omega) = false$

event $a \wedge b$ = points where $A(\omega) = true$ and $B(\omega) = true$

Often in AI applications, the sample points are **defined**

by the values of a set of random variables, i.e., the

sample space is the Cartesian product of the ranges of the variables

With Boolean variables, sample point = propositional logic model

e.g., $A = true$, $B = false$, or $a \wedge \neg b$.

Proposition = disjunction of atomic events in which it is true

$$\text{e.g., } (a \vee b) \equiv (\neg a \wedge \neg b) \vee (a \wedge \neg b) \vee (a \wedge b)$$


$$\Rightarrow P(a \vee b) = P(\neg a \wedge \neg b) + P(a \wedge \neg b) + P(a \wedge b)$$

Chapter 13 9

Why use probability?

The definitions imply that certain logically related events must have related probabilities

E.g., $P(a \vee b) = P(a) + P(b) - P(a \wedge b)$



de Finetti (1931): an agent who bets according to probabilities that violate these axioms can be forced to bet so as to lose money regardless of outcome.

Chapter 13 10

Syntax for propositions

Propositional or **Boolean** random variables

e.g., $Cavity$ (do I have a cavity?)

$Cavity = true$ is a proposition, also written $cavity$

Discrete random variables (**finite** or **infinite**)

e.g., $Weather$ is one of $\{sunny, rain, cloudy, snow\}$

$Weather = rain$ is a proposition

Values must be exhaustive and mutually exclusive

Continuous random variables (**bounded** or **unbounded**)

e.g., $Temp = 21.6$; also allow, e.g., $Temp < 22.0$.

Arbitrary Boolean combinations of basic propositions

Chapter 13 11

Prior probability

Prior or **unconditional** probabilities of propositions

e.g., $P(Cavity = true) = 0.1$ and $P(Weather = sunny) = 0.72$

correspond to belief prior to arrival of any (new) evidence

Probability distribution gives values for all possible assignments:

$P(Weather) = \langle 0.72, 0.1, 0.08, 0.1 \rangle$ (**normalized**, i.e., sums to 1)

Joint probability distribution for a set of r.v.s gives the

probability of every atomic event on those r.v.s (i.e., every sample point)

$P(Weather, Cavity) = a 4 \times 2$ matrix of values:

$Weather =$	$sunny$	$rain$	$cloudy$	$snow$
$Cavity = true$	0.144	0.02	0.016	0.02
$Cavity = false$	0.576	0.08	0.064	0.08


Every question about a domain can be answered by the joint distribution because every event is a sum of sample points

Chapter 13 12

Probability for continuous variables

Express distribution as a parameterized function of value:

$P(X=x) = U[18, 26](x)$ = uniform density between 18 and 26



Here P is a density; integrates to 1.


$P(X=20.5) = 0.125$ really means

$$\lim_{dx \rightarrow 0} P(20.5 \leq X \leq 20.5 + dx)/dx = 0.125$$

Chapter 13 13

Gaussian density

$$P(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-(x-\mu)^2/2\sigma^2}$$



Chapter 13 14

Conditional probability

Conditional or posterior probabilities

e.g., $P(\text{cavity}|\text{toothache}) = 0.8$

i.e., **given that toothache is all I know**

NOT "if toothache then 80% chance of cavity"

(Notation for conditional distributions:

$\mathbf{P}(\text{Cavity}|\text{Toothache})$ = 2-element vector of 2-element vectors)

If we know more, e.g., cavity is also given, then we have

$P(\text{cavity}|\text{toothache}, \text{cavity}) = 1$

Note: the less specific belief **remains valid** after more evidence arrives, but is not always **useful**

New evidence may be irrelevant, allowing simplification, e.g.,

$P(\text{cavity}|\text{toothache}, \text{9ersWin}) = P(\text{cavity}|\text{toothache}) = 0.8$

This kind of inference, sanctioned by domain knowledge, is crucial

Chapter 13 15

Conditional probability

Definition of conditional probability:

$$P(a|b) = \frac{P(a \wedge b)}{P(b)} \text{ if } P(b) \neq 0$$

Product rule gives an alternative formulation:

$$P(a \wedge b) = P(a|b)P(b) = P(b|a)P(a)$$

A general version holds for whole distributions, e.g.,

$$\mathbf{P}(\text{Weather}, \text{Cavity}) = \mathbf{P}(\text{Weather}|\text{Cavity})\mathbf{P}(\text{Cavity})$$

(View as a 4×2 set of equations, **not** matrix mult.)

Chain rule is derived by successive application of product rule:

$$\begin{aligned} P(X_1, \dots, X_n) &= P(X_1, \dots, X_{n-1}) P(X_n|X_1, \dots, X_{n-1}) \\ &= P(X_1, \dots, X_{n-2}) P(X_{n-1}|X_1, \dots, X_{n-2}) P(X_n|X_1, \dots, X_{n-1}) \\ &= \dots \\ &= \prod_{i=1}^n P(X_i|X_1, \dots, X_{i-1}) \end{aligned}$$

Chapter 13 16

Inference by enumeration

Start with the joint distribution:

	toothache		\neg toothache	
	catch	\neg catch	catch	\neg catch
cavity	.108	.012	.072	.008
\neg cavity	.016	.064	.144	.576

For any proposition ϕ , sum the atomic events where it is true:

$$P(\phi) = \sum_{\omega: \omega \models \phi} P(\omega)$$

Chapter 13 17

Inference by enumeration

Start with the joint distribution:

	toothache		\neg toothache	
	catch	\neg catch	catch	\neg catch
cavity	.108	.012	.072	.008
\neg cavity	.016	.064	.144	.576

For any proposition ϕ , sum the atomic events where it is true:

$$P(\phi) = \sum_{\omega: \omega \models \phi} P(\omega)$$

$$P(\text{toothache}) = 0.108 + 0.012 + 0.016 + 0.064 = 0.2$$

Chapter 13 18

Inference by enumeration

Start with the joint distribution:

	toothache		¬ toothache	
	catch	¬ catch	catch	¬ catch
cavity	.108	.012	.072	.008
¬ cavity	.016	.064	.144	.576

For any proposition ϕ , sum the atomic events where it is true:

$$P(\phi) = \sum_{\omega:\omega\models\phi} P(\omega)$$

$$P(\text{cavity} \vee \text{toothache}) = 0.108 + 0.012 + 0.072 + 0.008 + 0.016 + 0.064 = 0.28$$

Inference by enumeration

Start with the joint distribution:

	toothache		¬ toothache	
	catch	¬ catch	catch	¬ catch
cavity	.108	.012	.072	.008
¬ cavity	.016	.064	.144	.576

Can also compute conditional probabilities:

$$P(\neg \text{cavity} | \text{toothache}) = \frac{P(\neg \text{cavity} \wedge \text{toothache})}{P(\text{toothache})} = \frac{0.016 + 0.064}{0.108 + 0.012 + 0.016 + 0.064} = 0.4$$

Normalization

	toothache		¬ toothache	
	catch	¬ catch	catch	¬ catch
cavity	.108	.012	.072	.008
¬ cavity	.016	.064	.144	.576

Denominator can be viewed as a normalization constant α

$$\begin{aligned} P(\text{Cavity} | \text{toothache}) &= \alpha P(\text{Cavity}, \text{toothache}) \\ &= \alpha [P(\text{Cavity}, \text{toothache}, \text{catch}) + P(\text{Cavity}, \text{toothache}, \neg \text{catch})] \\ &= \alpha [(0.108, 0.016) + (0.012, 0.064)] \\ &= \alpha (0.12, 0.08) = (0.6, 0.4) \end{aligned}$$

General idea: compute distribution on query variable by fixing evidence variables and summing over hidden variables

Inference by enumeration, contd.

Let \mathbf{X} be all the variables. Typically, we want the posterior joint distribution of the query variables \mathbf{Y} given specific values \mathbf{e} for the evidence variables \mathbf{E}

Let the hidden variables be $\mathbf{H} = \mathbf{X} - \mathbf{Y} - \mathbf{E}$

Then the required summation of joint entries is done by summing out the hidden variables:

$$P(\mathbf{Y} | \mathbf{E} = \mathbf{e}) = \alpha P(\mathbf{Y}, \mathbf{E} = \mathbf{e}) = \alpha \sum_{\mathbf{h}} P(\mathbf{Y}, \mathbf{E} = \mathbf{e}, \mathbf{H} = \mathbf{h})$$

The terms in the summation are joint entries because \mathbf{Y} , \mathbf{E} , and \mathbf{H} together exhaust the set of random variables


Obvious problems:

- 1) Worst-case time complexity $O(d^n)$ where d is the largest arity
- 2) Space complexity $O(d^n)$ to store the joint distribution
- 3) How to find the numbers for $O(d^n)$ entries???

Independence

A and B are independent iff

$$P(A|B) = P(A) \quad \text{or} \quad P(B|A) = P(B) \quad \text{or} \quad P(A, B) = P(A)P(B)$$



$$P(\text{Toothache}, \text{Catch}, \text{Cavity}, \text{Weather}) = P(\text{Toothache}, \text{Catch}, \text{Cavity})P(\text{Weather})$$

32 entries reduced to 12; for n independent biased coins, $2^n \rightarrow n$

Absolute independence powerful but rare

Dentistry is a large field with hundreds of variables, none of which are independent. What to do?

Conditional independence

$P(\text{Toothache}, \text{Cavity}, \text{Catch})$ has $2^3 - 1 = 7$ independent entries

If I have a cavity, the probability that the probe catches in it doesn't depend on whether I have a toothache:

$$(1) P(\text{catch} | \text{toothache}, \text{cavity}) = P(\text{catch} | \text{cavity})$$

The same independence holds if I haven't got a cavity:

$$(2) P(\text{catch} | \text{toothache}, \neg \text{cavity}) = P(\text{catch} | \neg \text{cavity})$$

Catch is conditionally independent of Toothache given Cavity :

$$P(\text{Catch} | \text{Toothache}, \text{Cavity}) = P(\text{Catch} | \text{Cavity})$$

Equivalent statements:

$$P(\text{Toothache} | \text{Catch}, \text{Cavity}) = P(\text{Toothache} | \text{Cavity})$$

$$P(\text{Toothache}, \text{Catch} | \text{Cavity}) = P(\text{Toothache} | \text{Cavity})P(\text{Catch} | \text{Cavity})$$

Conditional independence contd.

Write out full joint distribution using chain rule:

$$\begin{aligned} & \mathbf{P}(\text{Toothache}, \text{Catch}, \text{Cavity}) \\ &= \mathbf{P}(\text{Toothache}|\text{Catch}, \text{Cavity})\mathbf{P}(\text{Catch}, \text{Cavity}) \\ &= \mathbf{P}(\text{Toothache}|\text{Catch}, \text{Cavity})\mathbf{P}(\text{Catch}|\text{Cavity})\mathbf{P}(\text{Cavity}) \\ &= \mathbf{P}(\text{Toothache}|\text{Cavity})\mathbf{P}(\text{Catch}|\text{Cavity})\mathbf{P}(\text{Cavity}) \end{aligned}$$

I.e., $2 + 2 + 1 = 5$ independent numbers (equations 1 and 2 remove 2)

In most cases, the use of conditional independence reduces the size of the representation of the joint distribution from exponential in n to linear in n .

Conditional independence is our most basic and robust form of knowledge about uncertain environments.

Bayes' Rule

Product rule $P(a \wedge b) = P(a|b)P(b) = P(b|a)P(a)$

$$\Rightarrow \text{Bayes' rule } P(a|b) = \frac{P(b|a)P(a)}{P(b)}$$

or in distribution form

$$\mathbf{P}(Y|X) = \frac{\mathbf{P}(X|Y)\mathbf{P}(Y)}{\mathbf{P}(X)} = \alpha\mathbf{P}(X|Y)\mathbf{P}(Y)$$

Useful for assessing diagnostic probability from causal probability:

$$P(\text{Cause}|\text{Effect}) = \frac{P(\text{Effect}|\text{Cause})P(\text{Cause})}{P(\text{Effect})}$$

E.g., let M be meningitis, S be stiff neck:

$$P(m|s) = \frac{P(s|m)P(m)}{P(s)} = \frac{0.8 \times 0.0001}{0.1} = 0.0008$$

Note: posterior probability of meningitis still very small!

Bayes' Rule and conditional independence

$$\begin{aligned} & \mathbf{P}(\text{Cavity}|\text{toothache} \wedge \text{catch}) \\ &= \alpha \mathbf{P}(\text{toothache} \wedge \text{catch}|\text{Cavity})\mathbf{P}(\text{Cavity}) \\ &= \alpha \mathbf{P}(\text{toothache}|\text{Cavity})\mathbf{P}(\text{catch}|\text{Cavity})\mathbf{P}(\text{Cavity}) \end{aligned}$$

This is an example of a naive Bayes model:

$$\mathbf{P}(\text{Cause}, \text{Effect}_1, \dots, \text{Effect}_n) = \mathbf{P}(\text{Cause})\prod_i \mathbf{P}(\text{Effect}_i|\text{Cause})$$



Total number of parameters is linear in n

Wumpus World

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 B OK	2,2	3,2	4,2
1,1 OK	2,1 B OK	3,1	4,1

$P_{ij} = \text{true}$ iff $[i, j]$ contains a pit

$B_{ij} = \text{true}$ iff $[i, j]$ is breezy

Include only $B_{1,1}, B_{1,2}, B_{2,1}$ in the probability model

Specifying the probability model

The full joint distribution is $\mathbf{P}(P_{1,1}, \dots, P_{4,4}, B_{1,1}, B_{1,2}, B_{2,1})$

Apply product rule: $\mathbf{P}(B_{1,1}, B_{1,2}, B_{2,1} | P_{1,1}, \dots, P_{4,4})\mathbf{P}(P_{1,1}, \dots, P_{4,4})$

(Do it this way to get $P(\text{Effect}|\text{Cause})$.)

First term: 1 if pits are adjacent to breezes, 0 otherwise

Second term: pits are placed randomly, probability 0.2 per square:

$$\mathbf{P}(P_{1,1}, \dots, P_{4,4}) = \prod_{i,j=1,1}^{4,4} \mathbf{P}(P_{i,j}) = 0.2^n \times 0.8^{16-n}$$

for n pits.

Observations and query

We know the following facts:

$$b = \neg b_{1,1} \wedge b_{1,2} \wedge b_{2,1}$$

$$\text{known} = \neg p_{1,1} \wedge \neg p_{1,2} \wedge \neg p_{2,1}$$

Query is $\mathbf{P}(P_{1,3}|\text{known}, b)$

Define $\text{Unknown} = P_{ij}$ s other than $P_{1,3}$ and Known


For inference by enumeration, we have

$$\mathbf{P}(P_{1,3}|\text{known}, b) = \alpha \sum_{\text{unknown}} \mathbf{P}(P_{1,3}, \text{unknown}, \text{known}, b)$$

Grows exponentially with number of squares!

Using conditional independence

Basic insight: observations are conditionally independent of other hidden squares given neighbouring hidden squares




Define $Unknown = Fringe \cup Other$
 $P(b|P_{1,3}, Known, Unknown) = P(b|P_{1,3}, Known, Fringe)$

Manipulate query into a form where we can use this!

Using conditional independence contd.

$$\begin{aligned}
 P(P_{1,3}|known, b) &= \alpha \sum_{unknown} P(P_{1,3}, unknown, known, b) \\
 &= \alpha \sum_{unknown} P(b|P_{1,3}, known, unknown) P(P_{1,3}, known, unknown) \\
 &= \alpha \sum_{fringe} \sum_{other} P(b|known, P_{1,3}, fringe, other) P(P_{1,3}, known, fringe, other) \\
 &= \alpha \sum_{fringe} \sum_{other} P(b|known, P_{1,3}, fringe) P(P_{1,3}, known, fringe, other) \\
 &= \alpha \sum_{fringe} P(b|known, P_{1,3}, fringe) \sum_{other} P(P_{1,3}, known, fringe, other) \\
 &= \alpha \sum_{fringe} P(b|known, P_{1,3}, fringe) \sum_{other} P(P_{1,3}) P(known) P(fringe) P(other) \\
 &= \alpha P(known) P(P_{1,3}) \sum_{fringe} P(b|known, P_{1,3}, fringe) P(fringe) \sum_{other} P(other) \\
 &= \alpha' P(P_{1,3}) \sum_{fringe} P(b|known, P_{1,3}, fringe) P(fringe)
 \end{aligned}$$

Using conditional independence contd.



$$P(P_{1,3}|known, b) = \alpha' \langle 0.2(0.04 + 0.16 + 0.16), 0.8(0.04 + 0.16) \rangle \approx \langle 0.31, 0.69 \rangle$$

$$P(P_{2,2}|known, b) \approx \langle 0.86, 0.14 \rangle$$

Summary

Probability is a rigorous formalism for uncertain knowledge

Joint probability distribution specifies probability of every atomic event

Queries can be answered by summing over atomic events

For nontrivial domains, we must find a way to reduce the joint size

Independence and conditional independence provide the tools