

# Subduction origin on early Earth: A hypothesis

Vicki L. Hansen

Department of Geological Sciences, University of Minnesota–Duluth, Duluth, Minnesota 55812, USA

## ABSTRACT

**I propose a hypothesis for the origin of subduction on early Earth that is directly coupled with mantle upwelling and the formation of mafic crust. The hypothesis invokes spatial and temporal overlap of both endogenic and exogenic processes, broad quasi-cylindrical mantle upwelling and large bolide impact, respectively, leading to subduction and spreading, two signature processes of modern plate tectonics. The spatial and temporal intersection of these processes could have occurred variably across early Earth, and thus subduction could have begun at different locations and times globally. The hypothesis postulates that the ability of a terrestrial planet to evolve plate tectonics results from a balance between the strength of its lithosphere and the size bolide it can attract and survive; both factors, to a first order, are a function of planet size.**

**Keywords:** subduction origin, impact, Archean, bolide, plate tectonics origin.

## INTRODUCTION

How and when plate tectonics began has been enthusiastically debated for decades. The debate currently combines a quest for the oldest evidence of plate tectonic processes (e.g., Cawood et al., 2006; Condie et al., 2006; Witze, 2006; Furnes et al., 2007). Modern plate tectonic processes embrace both the formation of mafic crust and the recycling of mafic crust to the mantle through subduction. Subduction is like a virus, once begun it can easily spread, yet a mechanism for inception of the subduction process has thus far proven elusive (e.g., Hilde et al., 1977; Casey and Dewey, 1984; Cloetingh et al., 1989; Mueller and Phillips, 1991). Thus, how subduction began is directly tied to the debate about the initiation of modern plate tectonic processes. As stated by Lin (2007), there must have been a time on Earth when plate tectonics did not operate and, thus, a process by which subduction began. I propose a hypothesis for the origin of subduction, coupled with the formation of mafic crust, that calls for serendipitous spatial intersection of endogenic processes (broad quasi-cylindrical mantle upwelling) and exogenic events (large bolide impacts) on early Earth. The hypothesis addresses the formation of long (>1000 km) fundamental boundaries, marked by contrasts in composition, rheology, and density, that evolve into subduction zones.

## EARLY EARTH

Early Earth differed from contemporary Earth in at least two major ways. (1) The planet was pummeled by extraterrestrial bolides, large and small; and (2) early Earth had significantly more thermal energy. The first is clear from solar system evolution models and impact craters and basins on the Moon and Mars. The second may have resulted in a stagnant-lid regime,

rather than a convective-lid regime (Tackley, 2000; Korenaga, 2006), hampering rather than enabling lithosphere-scale recycling.

Although there is much debate about the environmental conditions of early Earth, it is generally agreed that higher continental geothermal gradients, a result of greater radiogenic heat production and stronger basal heat flow, existed in the Archean (Taylor and McLennan, 1986; Bickle, 1978; Hoffman and Ranalli, 1988; Davies, 1992). An elevated geothermal gradient would lead to increased crustal buoyancy, increased ductility, and lower overall crustal strength; it would limit the ability of continental crust to develop sharp gradients in thickness or strength (Choukroune et al., 1995; Rey and Houseman, 2006). The broadly ductile nature of this crust would therefore inhibit the formation of abrupt compositional and/or rheological boundaries.


Fundamental, relatively sharp, focused boundaries form the essence of modern plate tectonics. Rheological and density differences play critical roles in the nature of tectonic processes of all terrestrial planets. Subduction ultimately requires plates—large tracts of crust or lithosphere, marked by differences in density and, possibly, strength. Lower plates, those to be subducted, must have both elastic strength and high density relative to the ductile upper mantle. In contrast, low-density upper plates need not have elastic strength, they can deform in either a ductile or brittle fashion. So perhaps we might ask, how could early Earth form at least local regions of high-strength crust, or relatively sharp boundaries marked by pronounced rheological and/or density contrast? Composition, temperature, strain rate, and fluids affect, to first order, the brittle (strong) versus ductile (weak) character of silicate crust; here I consider the first three

variables to propose a mechanism by which early Earth might form long (>1000 km), sharp, fundamental boundaries that separate large tracts of crust with contrasting density and/or rheology. The fourth variable, fluids, enters the discussion with an Earth-Venus comparison.

## HYPOTHESIS FOR THE ORIGIN OF SUBDUCTION

A series of time-step cartoons illustrates salient features of the hypothesis, beginning with differentiated (buoyant) crust rich in radiogenic heat-producing elements (Fig. 1). Part of the crust resides above a broad quasi-cylindrical mantle upwelling where the crust is thinned thermally and/or mechanically (Time 1). Mantle-scale processes modify the crust from below, resulting in two types of crust: unmodified and thinned.

While the mantle may modify the crust from below, extraterrestrial bolides can modify the crust from above. Bolides intersected the Earth as they traveled around the young Sun. Among the many objects to strike early Earth, a large bolide impacts the surface overlapping with the trace of already thinned felsic crust (Time 2). This impact event excavates a huge region of crust (>1000 km diameter), displacing much of it upward and outward, resulting in a vast region of a third type of crust that is thin, shocked, and weak (Time 3). The excavated crust, marking the impact site, could be expected to have a sharp boundary (strain gradient) against adjacent relatively unmodified (or now thickened) crust as a result of the extremely high strain rate experienced during the impact event. Although the crust would respond in a ductile fashion to low strain rate mantle flow from below, hypervelocity impact from above would induce a brittle response.



**Figure 1. Cartoon sequence (Time 1–Time 9; Time 5 repeated for clarity) illustrating the subduction origin hypothesis; schematic cross-sections A–B are shown for several panels. See text for discussion.**

The bolide, which penetrates the crust, also triggers a mantle response that in turn affects the surface (Time 4). Local superheating of the mantle causes massive partial melting, forming ultramafic to mafic melt that rises, forming new crust; impact could further spawn the formation of a relatively long lived mantle thermal anomaly (Jones et al., 2005; Elkins-Tanton and Hager, 2005). The region of excavated and thinned crust might be expected to respond much like thin lithosphere above a plume head, forming a ring-shaped trough that moves outward as the plume interacts with the thinned lithosphere (Griffiths and Campbell, 1991). The thermal

anomaly causes continued partial melting; melt rises to feed a nascent spreading center, which forms in the region inside the trough. As the trough moves outward, new mafic (to ultramafic) crust (a fourth type of crust) continues to form at the spreading center (Time 5). The broad mantle upwelling would influence the trend of the spreading center. Spreading could be symmetric or asymmetric depending on regional stress conditions; local buttressing by relatively immobile lithosphere might favor asymmetric spreading, rather than symmetrical spreading at modern ocean ridges. As new mafic crust is displaced outward, away from the spreading center,

the region of excavated and thinned felsic crust shortens and thickens. The nature of crustal thickening would depend on the inherent rheology of the excavated and thinned crust. One might reasonably predict a ductile response due to a high geothermal gradient, although the specific behavior is not critical to the hypothesis. What is important is that the excavated and thinned crust is progressively displaced outward with formation of new mafic crust. The spreading center propagates laterally in a direction parallel to the broad mantle upwelling; the growing tract of mafic crust progressively displaces the excavated and thinned crust until the new mafic crust reaches the limit of the original, impact-softened and excavated crust (Time 6).

Once the mafic crust reaches the spatial/rheological limit of the excavated crustal region, and given the right conditions (i.e., compositional and thermal density balanced against spreading rate and relative strength of mafic and felsic crusts), the early-formed mafic crust would begin to sink (Time 7), and then subduct, beneath the buoyant crust, which would become the upper plate (Time 8). The spatial/rheological limit refers to the point at which the felsic crust (modified or not) begins to resist displacement, becoming a stress concentrator (Cloetingh et al., 1989), caused by further generation of mafic crust at the spreading center. The spreading center continues to propagate along strike akin to modern ridges, and it could ultimately penetrate the felsic crust left unmodified by the impact event, but where it was thinned above the original mantle upwelling. As the system evolves, transform faults might form as dictated by geometric and kinematic criteria along portions of the mafic-felsic crust boundary. The subduction zone could propagate along its length, as did the spreading center (Time 9). With the concurrent formation of new mafic crust and subsequent subduction of older mafic crust, plate tectonics have begun in this region of the Earth.

This hypothesis calls upon serendipitous spatial overlap of both endogenic and exogenic processes. If a large bolide fell on crust far removed from a mantle upwelling, the events proposed here might not evolve into mafic crust formation and recycling via subduction, because there would be no upwelling to act as a stress guide for crustal extension. Furthermore, because the spreading center might not be able to propagate into unmodified crust, the overall areal extent of newly formed mafic crust would be limited. That is, unmodified crust might resist spreading center propagation. The proposed process could become arrested at any stage described; in this case, subduction would not begin.

The hypothesis, which assumes a constant radius Earth, accommodates, and essentially requires, concurrent formation and recycling of mafic crust. The inception of subduction requires

formation of a sufficiently large tract of mafic crust to allow for crustal cooling, and this crust must form a strong (i.e., elastic) brittle plate of significant density difference compared to the asthenosphere, but also compared to the upper plate, such that the newly formed mafic crust will be thrust below the felsic crust. The size of the impact basin is important in that an extensive tract of mafic crust must form to allow subduction to begin. The question of whether bolides of sufficient size existed is evident by examination of Earth's neighbors, Mars and the Moon, which preserve at least partial records of that tumultuous time. Mars, about half the diameter of Earth and 11% of its mass, preserves several impact basins with diameters >1800 km (Argyre, 1800 km; Isidis, 1900 km; Hellus, 2300 km; Utopia, 3000 km). The Moon, less than one-third Earth's diameter and 1.2% its mass, also has huge impact basins (Orientale, 930 km; Imbrium, 1160 km; South Pole–Aitken, 2300 km). Earth, with significantly greater mass, should have sustained similarly large impact events as these smaller neighbors, forming impact basins of at least equal size. Subduction zones associated with the Lesser Antilles and South Sandwich Islands, with diameters of ~1000 km, might be taken as a minimum size for impact basins that might evolve into spreading-subduction processes.

### CONSEQUENCES AND RESERVATIONS

Given that such large impact basins occur on Mars and the Moon, one might ask why plate tectonics did not begin on these bodies. The hypothesis presented here embraces a balance between a terrestrial body's lithospheric strength and bolide size. A bolide must form a huge impact basin and penetrate the lithosphere. If a bolide is too small, or the lithosphere is too strong, plate tectonics will not occur. To a first order, lithospheric strength for terrestrial bodies is a function of planet size, or its surface area:volume ratio,  $A/V$  (Solomon and Head, 1982). Small bodies, with high  $A/V$ , have strong lithospheres. With a lower  $A/V$ , a body's heat budget increases, and its lithosphere is weaker. The size bolide a planet can attract and sustain is also a direct function of mass, and therefore size.

Thus, plate tectonics (subduction-spreading processes) might only form on large solid-surfaced planets—bodies with relatively weak lithospheres that can sustain the impact of very large bolides. In effect, there could be a minimum size at which a planet might develop plate tectonics. The Moon is too small. Any bolide that the Moon could sustain would not initiate subduction-spreading processes because of its thick lithosphere. Mars also has a thick lithosphere, although this has not always been the case. Early in Mars' history it had significantly more heat and, as a result, a thinner and weaker lithosphere. At that same time the solar system

contained numerous large meteoroids. A single large bolide (Wilhelms and Squyres, 1984) or multiple large bolides (McGill, 1989; Frey and Schultz, 1988) may have collided with what is currently Mars' Northern Hemisphere, forming Mars' crustal dichotomy. The impact event(s) could have, in turn, spawned an endogenic response to develop a spreading center forming mafic crust, which was displaced asymmetrically outward and subsequently subducted along a portion of the initial impact boundary (Sleep, 1994). Thus the formation of Mars' crustal dichotomy might have required partnership between exogenic and endogenic events (Wise et al., 1979; McGill and Dimitriou, 1990; Zuber, 2001), which evolved into plate tectonic processes (Sleep, 1994). As Mars cooled, due mainly to its high  $A/V$  but enhanced by nascent mafic crust formation, plate tectonic processes would have halted abruptly, almost as quickly as they began, preserving a truly ancient ocean-like basin across Mars' Northern Hemisphere (Frey, 2006).

Venus, with ~80% of Earth's mass, lacks plate tectonics. Thus, (1) Venus might mark a lower limit for the terrestrial planet mass required for plate tectonics to evolve; or (2) Venus once had plate tectonics, but fast cooling led to its demise; or (3) Venus's dry strong crust (Mackwell et al., 1998) rendered its lithosphere too strong for plate tectonics to develop. Scenario 1 seems unlikely given the similarity in Earth and Venus size and composition. Scenario 2 is also unlikely given the role of a dense atmosphere in maintaining a hot interior (e.g., Abe and Matsui, 1988), and given that Venus's supercritical  $\text{CO}_2$ -rich atmosphere acts more like a conducting layer than a convective layer with regard to heat transfer (Snyder, 2002). The third option highlights lithospheric rheology, a fundamental component of the bolide model. Water plays a critical role in subduction initiation (and hence plate tectonics) due to the mode of failure of silicate lithospheres (Regenauer-Lieb et al., 2001). Convergence across a dry lithosphere (such as Venus) results in focused brittle failure of the upper lithosphere, while the bottom of the lithosphere deforms in a diffuse fashion and delaminates and recycles to the mantle, leaving the surface crust relatively intact. In contrast, wet lithosphere fails across its entire mechanical thickness, resulting in the formation of a narrow fault-like zone through the lithosphere; in this case the entire lithospheric thickness recycles to the mantle (Regenauer-Lieb et al., 2001). Once initiated, subduction of the lithosphere provides a means of recycling water to the mantle, triggering a double feedback mechanism (thermoelastic and thermal-rheological) to promote plate tectonics (Regenauer-Lieb et al., 2001). Thus, Venus's lack of plate tectonics might highlight the role of rheology across the entire thickness of a planet's lithosphere.

Inception of plate tectonics by the proposed hypothesis would not require strictly serendipitous occurrence of exogenic and endogenic processes. Rather, initiation of plate tectonics might be a function of planet size, which in turn influences both endogenic (heat budget, cooling rate, and lithospheric strength) and exogenic (bolide attractor) factors in its formative years. Water also plays a critical role in plate tectonic process initiation and continued evolution and should be considered in cases where  $A/V$  criteria are met. Plate tectonic initiation by the mechanism proposed here is also limited by the availability of large bolides, and calls for consideration of the thermomechanical evolution of a planet within the context of solar system evolution. That is, the initiation of plate tectonic processes might require intersection of temporal and spatial factors with regard to planet differentiation and cooling, and how these processes affect lithospheric strength, as well as solar system evolution, with particular regard to large bolides.

The effect of large bolides on an early Earth might provide clues to some unsolved mysteries of the Archean (Condie and Benn, 2006). For example, (1) bolide-induced initiation of subduction spreading by the mechanism suggested here could allow for two very different styles of crustal deformation—diapirism-sagduction and plate tectonics—to operate concurrently in the Archean. (2) Komatiites, which characterize the Archean, could result from superheating of the mantle due to large bolide impact (high-temperature komatiite), or heating and hydration (hydrous komatiite) due to icy bolide impact. In either case, (3) the resulting mantle residuum (Jordan, 1978) could, in turn, form low-density keels of Archean lithosphere, the formation of which is controversial. Komatiites would be rapidly recycled, whereas the restite—buoyant, refractory and cool—would survive. (4) Given that thermal modeling indicates that mantle-formed plumes might have been weak to nonexistent in the Archean (Korenaga, 2006), perhaps bolides (or bolide-spawned plumes) triggered the ca. 2.75 Ga global events credited to plumes. In addition, if bolides played a critical role in the origin of plate tectonics, then with plate tectonics came crucial extraterrestrial ingredients for life on Earth, delivered to a favorable environment that literally nursed them to life.

One might argue that calling on impact processes violates uniformitarianism, and thus the very roots of geology. However, perhaps ignoring impact events represents a modern form of geocentric provincialism. Both secular and uniformitarian philosophies, each temporal- and spatial-dependent approaches, are critical to understanding Earth's formation and evolution. Early Earth differs from contemporary Earth; today's catastrophe could have been yesterday's norm. Recent work indicates that Earth's

differentiation occurred earlier than previously thought (Boyet and Carlson, 2005), yet major impact events (e.g., late heavy bombardment) may have continued over a broader time span, including 4.1–2.0 Ga (Norman et al., 2006; Hartmann et al., 2007), than previously appreciated. Thus, there is every reason to suppose that large bolides did have strong effects on early Earth. A wide range of petrologic, rheological, thermal, and/or heat flow arguments might be constructed to identify strengths, weakness, or fundamental flaws with the proposed hypothesis, which outlines one possible way that bolides might have played a critical role in Earth's evolution.

#### ACKNOWLEDGMENTS

I thank P. Cawood, P. Schultz, and an anonymous reviewer for careful and insightful reviews, and J. Goodge and J. Swenson for comments on earlier versions of the manuscript. I thank the McKnight Foundation and the National Aeronautics and Space Administration (grants NNX06AB90G and NNG05GM26G) for support.

#### REFERENCES CITED

- Abe, Y., and Matsui, T., 1988, Evolution of an impact-generated H<sub>2</sub>O-CO<sub>2</sub> atmosphere and formation of a hot proto-ocean on Earth: *Journal of the Atmospheric Sciences*, v. 45, p. 3081–3101, doi: 10.1175/1520-0469(1988)045<3081:EOAIGH>2.0.CO;2.
- Bickle, M.J., 1978, Heat loss from the Earth: A constraint of Archean tectonics from the relation between geothermal gradients and the rate of plate production: *Earth and Planetary Science Letters*, v. 40, p. 301–315, doi: 10.1016/0012-821X(78)90155-3.
- Boyet, M., and Carlson, R.W., 2005, Nd-142 evidence for early (> 4.53 Ga) global differentiation of the silicate Earth: *Science*, v. 309, p. 576–581, doi: 10.1126/science.1113634.
- Casey, J.F., and Dewey, J.F., 1984, Initiation of subduction zones along transform and accreting plate boundaries, triple-junction evolution, and forearc spreading centers—Implications for ophiolitic geology and obduction, *in* Gass, I.G., et al., eds., *Ophiolites and oceanic lithosphere*: Geological Society [London] Special Publication 13, p. 269–290, doi: 10.1144/GSL.SP.1984.013.01.22.
- Cawood, P.A., Kroner, A., and Pisarevsky, S., 2006, Precambrian plate tectonics: Criteria and evidence: *GSA Today*, v. 16, no. 7, p. 4–11, doi: 10.1130/GSAT01607.1.
- Choukroune, P., Bouhallier, H., and Arndt, N.T., 1995, Soft lithosphere during periods of Archean crustal growth or crustal reworking, *in* Coward, M.P., and Ries, A.C., eds., *Early Precambrian processes*: Geological Society [London] Special Publication 95, p. 67–86.
- Cloetingh, S., Wortel, R., and Vlaar, N.J., 1989, On the initiation of subduction zones: *Pure and Applied Geophysics*, v. 129, p. 7–25, doi: 10.1007/BF00874622.
- Condie, K.C., and Benn, K., 2006, Archean geodynamics: Similar to or different from modern geodynamics?, *in* Benn, K., et al., eds., *Archean geodynamics and environments*: American Geophysical Union Geophysical Monograph 164, p. 47–59.
- Condie, K.C., Kröner, A., and Stern, R.J., 2006, When did plate tectonics begin?: *GSA Today*, v. 16, no. 10, p. 40–41, doi: 10.1130/1052-5173(2006)16[40:PCRWDP]2.0.CO;2.
- Davies, G.F., 1992, On the emergence of plate tectonics: *Geology*, v. 20, p. 963–966, doi: 10.1130/0091-7613(1992)020<0963:OTEOPT>2.3.CO;2.
- Elkins-Tanton, L., and Hager, B., 2005, Giant meteoroid impacts can cause volcanism: *Earth and Planetary Science Letters*, v. 239, p. 219–232, doi: 10.1016/j.epsl.2005.07.029.
- Frey, H.V., 2006, Impact constraints on, and a chronology for, major events in early Mars history: *Journal of Geophysical Research*, v. 111, p. E08S91, doi: 10.1029/2005JE002449.
- Frey, H., and Schultz, R.A., 1988, Large impact basins and the mega-impact origin for the crustal dichotomy on Mars: *Geophysical Research Letters*, v. 15, p. 229–232.
- Furnes, H., de Wit, M., Staudigel, H., Rosing, M., and Muehlenbachs, K., 2007, A vestige of Earth's oldest ophiolite: *Science*, v. 315, p. 1704–1707, doi: 10.1126/science.1139170.
- Griffiths, R.W., and Campbell, I.H., 1991, Interaction of mantle plume heads with the Earth's surface and onset of small-scale convection: *Journal of Geophysical Research*, v. 96, p. 18,295–18,310.
- Hartmann, W.K., Quantin, C., and Mangold, N., 2007, Possible long-term decline in impact rates—2. Lunar impact-melt data regarding impact history: *Icarus*, v. 186, p. 11–23, doi: 10.1016/j.icarus.2006.09.009.
- Hilde, T.W.C., Uyeda, S., and Kroenke, L., 1977, Evolution of the Western Pacific and its margin: *Tectonophysics*, v. 38, p. 145–165, doi: 10.1016/0040-1951(77)90205-0.
- Hoffman, P.F., and Ranalli, G., 1988, Archean oceanic flake tectonics: *Geophysical Research Letters*, v. 15, p. 1077–1080.
- Jones, A.P., Wunemann, K., and Price, D., 2005, Impact volcanism as a possible origin for the Ontong Java Plateau (OJP), *in* Foulger, G.R., et al., eds., *Plates, plumes, and paradigms*: Geological Society of America Special Paper 388, p. 711–720.
- Jordan, T.H., 1978, Composition and development of the continental tectosphere: *Nature*, v. 274, p. 544–548, doi: 10.1038/274544a0.
- Korenaga, J., 2006, Archean geodynamics and the thermal evolution of Earth, *in* Benn, K., et al., eds., *Archean geodynamics and environments*: American Geophysical Union Geophysical Monograph 164, p. 7–32.
- Lin, S., 2007, When did plate tectonics begin?: *GSA Today*, v. 17, no. 3, p. 12, doi: 10.1130/GSAT01703C.1.
- Mackwell, S.J., Zimmerman, M.E., and Kohlstedt, D.L., 1998, High-temperature deformation of dry diabase with application to tectonics on Venus: *Journal of Geophysical Research*, v. 103, p. 975–984, doi: 10.1029/97JB02671.
- McGill, G.E., 1989, Buried topography of Utopia, Mars: Persistence of a giant impact depression: *Journal of Geophysical Research*, v. 94, p. 2753–2759.
- McGill, G.E., and Dimitriou, A.M., 1990, Origin of the Martian global dichotomy by crustal thinning in the late Noachian or early Hesperian: *Journal of Geophysical Research*, v. 95, p. 12,595–12,605.
- Mueller, S., and Phillips, R.J., 1991, On the initiation of subduction: *Journal of Geophysical Research*, v. 96, p. 651–666.
- Norman, M.D., Duncan, R.A., and Huard, J.J., 2006, Identifying impact events within the lunar cataclysm from Ar-40-Ar-39 ages and compositions of Apollo 16 impact melt rocks: *Geochimica et Cosmochimica Acta*, v. 70, p. 6032–6049, doi: 10.1016/j.gca.2006.05.021.
- Regenauer-Lieb, K., Yuen, D.A., and Branlund, J., 2001, The initiation of subduction: Criticality by addition of water?: *Science*, v. 294, p. 578–580, doi: 10.1126/science.1063891.
- Rey, P.F., and Houseman, G., 2006, Lithospheric scale gravitational flow: The impact of body forces on orogenic processes from Archean to Phanerozoic, *in* Buitter, S.J.H., and Schreurs, G., eds., *Analogue and numerical modeling of crustal-scale processes*: Geological Society [London] Special Publication 253, p. 153–167.
- Sleep, N.H., 1994, Martian plate tectonics: *Journal of Geophysical Research*, v. 99, p. 5639–5655, doi: 10.1029/94JE00216.
- Snyder, D., 2002, Cooling of lava flows on Venus: The coupling of radiative and convective heat transfer: *Journal of Geophysical Research*, v. 107, no. E10, p. 5080–5088, doi: 10.1029/2001JE001501.
- Solomon, S.C., and Head, J.W., 1982, Mechanisms of lithospheric heat transport on Venus: Implications for tectonic style and volcanism: *Journal of Geophysical Research*, v. 87, p. 9236–9246.
- Tackley, P.J., 2000, Mantle convection and plate tectonics: Toward an integrated physical and chemical theory: *Science*, v. 288, p. 2002–2007, doi: 10.1126/science.288.5473.2002.
- Taylor, S.R., and McLennan, M.S., 1986, The chemical composition of the Archean crust, *in* Dawson, J.B., et al., eds., *The nature of the lower continental crust*: Geological Society [London] Special Publication 24, p. 173–178.
- Wilhelms, D.E., and Squyres, S.W., 1984, The Martian hemispheric dichotomy may be due to a giant impact: *Nature*, v. 309, p. 138–140, doi: 10.1038/309138a0.
- Wise, D.U., Golombek, M.P., and McGill, G.E., 1979, Tectonic evolution of Mars: *Journal of Geophysical Research*, v. 84, p. 7934–7939.
- Witze, A., 2006, The start of the world as we know it: *Nature*, v. 442, p. 128–131, doi: 10.1038/442128a.
- Zuber, M.T., 2001, The crust and mantle of Mars: *Nature*, v. 412, p. 220–227, doi: 10.1038/35084163.

Manuscript received 29 March 2007  
 Revised manuscript received 21 June 2007  
 Manuscript accepted 15 July 2007

Printed in USA