

SOUTHWEST

1000 feet=305 meters

SOUTHWEST

1000 feet=305 mete

NORTHWEST

1000 feet=305 meters

LAKE HAVASU CITY

MYLONITIC GNEISSES

MYLONITIC GNEISSES


**BOULDER MINE AREA** 


AUBREY HILLS

Large fault mullions and grooves (John, 1987a)

Note: Thin Quaternary units may be omitted

or exaggerated in cross sections.





K-Ar Biotite

5.9' Amphibolite K-Ar Hornblende

12 percent dikes

Seismic velocity boundar interpreted as the base of

highly faulted allochthons

K-Ar Hornblende

597+11

104±3

116±3

inferred from

models of gravity

gradient-denser rocks to the SW

(Simpson and

others, 1986)

1372±41 \*

Xap P81MH-10

8 percent dikes

Tfls Tvu Ysqm Qs<sub>2</sub>b Qs<sub>2</sub>a Tys Tfl Ysg Qs<sub>3</sub> Tys Qs<sub>3</sub> Tfg Qs<sub>2</sub> Qs<sub>4</sub> Ysg

43 Xga P81BK-13

44 Xga H83MH-67

CROSSMAN BLOCK

30 percent dikes

DETACHMENT

MOUNTAINS

STANDARD WASH AREA

MOHAVE

FAULT

MOUNTAINS DETACHMENT

MYLONITIC GNEISSES

45 **Xgl** H87MH-30

38.4'


41.8'

33.5'

8.9' Amphibolite

33.2' 12.6' Leucogneiss K-Ar Biotite

# ▲ diorite yenogranite Monzogranite



○ Tgd


□ TKg


▲ Kbg

TKqp 1

■ TKd 1

▲ XgI 3





average of 1000 points counted on stained slabs. International

MYLONITIC GNEISSES

NORTHEAST

Union of Geological Sciences classification (Streckeison, 1973).

# INTRODUCTION

- The Mohave Mountains area surrounds Lake Havasu City, Arizona, in the Basin and Range physiographic province (fig. 1). The Mohave Mountains and the Aubrey Hills form two northwest-trending ranges adjacent to Lake Havasu (elevation 132 m; 448 ft) on the Colorado River (fig. 2). The low Buck Mountains lie northeast of the Mohave Mountains in the alluviated valley of Dutch Flat. Lowlands at Standard Wash separate the Mohave Mountains from the Bill Williams Mountains to the southeast. The highest point in the area is Crossman Peak in the Mohave Mountains, at an elevation of 1519 m (5148 ft). Arizona Highway 95 is now rerouted in the northwestern part of the map area from its position portrayed on the base map; it now also passes through the southern edge of the map area (fig. 2).
- Geologic mapping was begun in 1980 as part of a program to assess the mineral resource potential of Federal lands under the jurisdiction of the U.S. Bureau of Land Management (Light and others, 1983). Mapping responsibilities were as follows: Proterozoic and Mesozoic rocks, K.A. Howard; dikes, J.K. Nakata; Miocene section, J.E. Nielson; and surficial deposits, H.G.
- Earlier geologic mapping includes reconnaissance mapping by Wilson and Moore (1959). The present series of investigations has resulted in reports on the crystalline rocks and structure (Howard and others, 1982a), dikes (Nakata, 1982), Tertiary stratigraphy (Pike and Hansen, 1982; Nielson, 1986; Nielson and Beratan, 1990), surficial deposits (Wilshire and Reneau, 1992), tectonics (Howard and John, 1987; Beratan and others, 1990), geophysics (Simpson and others, 1986), mineralization (Light and McDonnell, 1983; Light and others, 1983), field guides (Nielson, 1986; Howard and others, 1987), and geochronology (Nakata and others, 1990; Foster and others, 1990).
- The area lies within a terrane characterized by major low-angle normal faults (commonly called detachment faults) of Miocene age (Davis and others, 1980; Howard and John, 1987; John, 1987a: Davis and Lister, 1988: Davis, 1988: Carr, 1991). The Whipple Mountains detachment fault is exposed 3 km (2 mi) southwest of the mapped area, and the probably equivalent Chemehuevi detachment fault is exposed 8 km (5 mi) northwest of the mapped area. Both dip gently toward the Mohave Mountains area and project beneath it at shallow
- Proterozoic metamorphic and igneous rocks, Cretaceous plutons, and Tertiary dikes underlie the map area. Tertiary volcanic and sedimentary rocks overlie the Proterozoic rocks. Tilting to the southwest in the Miocene along northeast-dipping normal faults resulted in steep to overturned dips of lower Miocene strata and the Proterozoic-Tertiary nonconformity. Consequently the map view restores to cross-sectional views before the tilting, in which deeper crustal levels in any tilted block are encountered to the northeast. The faulting is inferred to relate to down-to-northeast motion on the deeper Whipple Mountains-Chemehuevi fault (Howard and
- John, 1987). Upper Miocene to Quaternary deposits overlie the deformed rocks. The largest of the tilted fault blocks is the Crossman block, centered in the map area (Crossman plate of Howard and others, 1982a). Other large tilt blocks are the Bill Williams, Buck, and Tumarion blocks (fig 3). Smaller tilt blocks or slices are present in the Standard Wash, Aubrey Hills, and Boulder Mine areas.
- Proterozoic and Mesozoic Rocks Granite gneisses are the predominant Early Proterozoic rocks in the map area. The gneisses are like those widely exposed elsewhere in the eastern Mojave Desert region (Wooden and Miller, 1990) and consist partly of supracrustal rocks and largely of metaplutonic rocks (Howard and others, 1982a). Scattered amphibolite bodies (Xga) help to define large folds in the gneiss terrane. Fold axes plunge to the southwest and restore to northeast plunge if Terti-
- ary tilting is removed (Howard and others, 1982a). Amphibolite is associated with white garnet-bearing gneissic pegmatite (Xgp) in the northwestern part of the Crossman block; it serves as a marker for fault separation to a block in the Boulder Mine area. The same assemblage also crops out along strike to the northeast in the Buck block, as well as farther southeast in the Crossman block.
- Remnant metasedimentary rocks (Xgs) and fine-grained biotite granite gneiss (Xgb) and younger foliated monzogranite (Xgp) crop out in the northern part of the area in structurally high positions of the pre-Tertiary crustal section. Augen gneiss (Xap, Xag, and Xagd) occurs in structurally intermediate levels of the section and becomes more foliated and more mafic with paleodepth. Dating by the uranium-lead (U-Pb) method indicates an Early Proterozoic age (1642 Ma) for an augen gneiss body in the Bill Williams Mountains (Wooden and Miller. 1990: \*table 1, No. 36). Howard and others (1982a) earlier had grouped the augen gneiss with Middle Proteozoic(?) granitoids of Standard Wash, which it resembles. The augen gneiss commonly coincides with aeromagnetic highs (Simpson and others, 1986). The deepest rocks in the exposed crustal section are dominated by garnet-spotted leucocratic gneiss (XgI) in which per-
- vasive garnets, now mostly retrograded, suggest high-grade metamorphism. Granite and quartz monzonite in Standard Wash (Ysg, Ysgm) are tentatively correlated with rocks of Middle Proterozoic age (about 1400 Ma) described by Anderson and Bender (1989). An aeromagnetic high is associated with the quartz monzodiorite (Simpson and others, 1986). The granite is seen to intrude older gneiss just south of the map area in the southern Aubrey Hills. The Standard Wash granitoids are not mapped in the Crossman, Bill Williams, or Buck blocks.
- Swarms of Middle Proterozoic ophitic diabase dikes intrude the gneiss and granitoids (see cross sections; Nakata, 1982; Howard, 1991). The dikes (Yd) are similar to diabase dikes distributed widely in Arizona and southeastern California and are interpreted to have an age of about 1100-1200 Ma (Wrucke and Shride, 1972; Hendricks and Lucchitta, 1974; Davis and others, 1980; Howard and others, 1982a; 1982b; Fitzgibbon, 1988). The dikes cut older folds and fabrics without deflection. Their generally northwest strikes and steep dips restore to subhorizontal when Tertiary tilting defined by the dip of lowest Tertiary strata is removed. Hence the dikes are interpreted as originally horizontal sheets. They are mapped to paleodepths of as great as 14 km below the Proterozoic-Tertiary nonconformity. Original depths of emplacement below the Proterozoic ground surface must have been even greater.
- Cooling ages obtained on Proterozoic rocks by the fission-track (zircon) and potassiumargon (K-Ar) methods range from 46 to 1372 Ma (table 1; Nakata and others, 1990). The numerical ages tend to young downward (northeastward) in the pre-Tertiary structural sections and are consistent with younger cooling at deeper levels. Age determinations by the  $^{40}$ Ar/ $^{39}$ Ar technique support this interpretation (Foster and others, 1990). Mineralized quartz veins in the Mohave Mountains contain sulfides and precious metals (Light and others, 1983). The veins cut the Proterozoic diabase and are cut by Tertiary dikes and have a minimum age of mid-Cretaceous as determined by isotopic dating: medium-grained white mica in altered gneiss associated with the veins yielded K-Ar ages of 90 to 102 Ma;
- numerical cooling ages on fine-grained white mica and zircon are 52 to 78 Ma (table 1, Nos. 30-32; Nakata and others, 1990). The Mohave Mountains and adjacent ranges are near the northeastern limit of the Cretaceous magmatic arc (Burchfiel and Davis, 1981; John, 1981; John and Wooden, 1990). Granitic plutons of known and inferred Cretaceous age are widespread to the west but are rare to the northeast. Cretaceous rocks in the Mohave Mountains area may be apophyses of the deeper batholithic terrane exposed to the west in the Chemehuevi Mountains (John, 1987b). This interpretation is based on a hypothesis of Howard and John (1987) that the northern Mohave Mountains represent the displaced upper plate to rocks exposed in the Chemehuevi
- area are correlated to Late Cretaceous rocks of the Chemehuevi Mountains Plutonic Suite (John, 1987b, 1988; John and Mukasa, 1990; John and Wooden, 1990). Rocks in the Mohave Mountains area that may be either Cretaceous or Tertiary in age include a northeast-striking rhyolite dike dated by K-Ar on biotite at 62 Ma (table 1, No. 28, unit TYd), northeast-striking dikes of lamprophyre and quartz porphyry (TKI, TKqp), and small stocks of granite and diorite (TKg, TKd) that yielded middle Tertiary numerical K-Ar ages on biotite and hornblende (table 1, Nos. 26, 27). Northeast-striking dikes of Laramide (Late Cretaceous to early Tertiary) age are widespread in Arizona (Rehrig and Heidrick, 1976).

Mountains. Granodiorite and porphyritic granite (Kgb, Kpg) in the northern part of the map

- Cenozoic Rocks The lowest stratified Tertiary unit (Tac) is arkosic conglomerate and sandstone, locally redbed, which is assigned an Oligocene or early Miocene age. It nonconformably overlies deeply
- oxidized Proterozoic rocks and forms the base of a heterogeneous, dominantly volcanic section \*Some ages listed in table 1 record cooling younger than age of rock (especially for Proterozoic rocks); other ages are

NORTHEAST

DETACHMENT

anomalously older than probable rock age (especially for dikes) and may result from excess argon (Nakata and others,

1990). Description of Map Units mentions ages that are believed to reflect crystallization age of sampled rock units.

BUCK MOUNTAINS

BUCK BLOCK

MOUNTAINS

Mapping of the Mohave Mountains area was a team effort. J.S. Noller, K.K. Beratan, D.M. Peterson, and D.V. Prose mapped parts of the area. R.D. Knox and M.A. Pernokas participated in sampling and field and laboratory study. We benefited from discussions with R.W. Simpson, W.A. Rehrig, T.D. Light, G.A. Davis, and J.L. Anderson. Critical reviews by D.R.

- Anderson, J.L., 1983, Proterozoic anorogenic granite plutonism of North America, in Meda-America Memoir 161, p. 133–154.
- Anderson, J.L., Barth, A.P., and Young, E.D., 1988, Mid-crustal roots of Cordilleran metamorphic core complexes: Geology, v. 16, p. 366–369. Anderson, J.L., and Rowley, M.C., 1981, Synkinematic intrusion of peraluminous and associ-
- Beratan, K.K., Blom, R.G., Nielson, J.E., and Crippen, R.E., 1990, Use of Landsat thematic mapper images in regional correlation of syntectonic strata, Colorado River extensional corridor, California and Arizona: Journal of Geophysical Research, v. 95, p. 615–624. Bull, W.B., 1974, Geomorphic tectonic analysis of the Vidal region, in Information concerning geology and seismology: Vidal Nuclear Generation Station, Vol. III, Appendix 2.5B.
- Geotectonic Development of California, Rubey Volume I: Englewood Cliffs, N.J., Prentice-Hall, Inc., p. 217–252.
- and Arizona: U.S. Geological Survey Professional Paper 1430, 40 p. Carr. W.J., Dickey, D.D., and Quinlivan, W.D., 1980, Geologic map of the Vidal NW, Vidal Junction, and parts of the Savahia Peak SW and Savahia Peak quadrangles, San Bernardino County, California: U.S. Geological Survey Miscellaneous Investigations Series Map

BILL WILLIAMS MOUNTAINS

FAULT

INTERIOR—GEOLOGICAL SURVEY, RESTON, VA-1999

- of otherwise early and middle Miocene age (fig. 4; Pike and Hansen, 1982; Nielson, 1986; Nielson and Beratan, 1990; Beratan and others, 1990). The volcanic rocks consist of mafic and silicic lava flows, volcaniclastic flows and breccias, air-fall tuff, and tuff breccia. The Peach Springs Tuff of Young and Brennan (1974), a regional ash-flow tuff, crops out high in the lower Miocene part of the section. The sedimentary rocks include arkosic conglomerate and conglomeratic sandstones derived from Proterozoic sources, sandstone and sedimentary breccia derived solely from reworked volcanic rocks, and fanglomerate, sandstone, and claystone from mixed sources. The lower part of the Tertiary section is intruded by silicic and mafic dikes that
- may be feeders for some of the flows. The dikes are part of a dense swarm of northeast-dipping Tertiary dikes that pervades the pre-Tertiary basement of the Crossman block. This Mohave Mountains dike swarm forms about 15 percent of the rock volume in the Crossman block (Nakata, 1982) and accounts for a northeast-southwest crustal dilation of 2-3 km that occurred during regional tectonic extension. The K-Ar ages considered most reliable as intrusive ages for the swarm range from 19.2±0.5 to

21.5±0.5 Ma (table 1, Nos. 21–25).

direction as older steeper tilts.

- The Tertiary strata below the Peach Springs Tuff correlate broadly with the sedimentary and volcanic rocks of Fox Wash described by Sherrod (1988) from the Bill Williams Mountains. Age determinations by the K-Ar method indicate that the age of the volcanic rocks in this section is about 19 to 22 Ma (table 1, Nos. 8-11, 13; Nielson and Beratan, 1990; Nakata and others, 1990). The Peach Springs Tuff is dated from Kingman, Arizona, at 18.5+0.2 Ma (Nielson and others, 1990). The steeply dipping to overturned oldest part of the Tertiary section is overlain succesively by units of early Miocene to middle Miocene age having progressively shallower dips, in places without obvious unconformities between strata. Therefore the stratified units were deposited during tilting, and they likely record sedimentary influxes that relate to the tilting and faulting. Conglomerates low in the section are largely roundstone. Higher units include lacustrine strata deposited in closed structural basins. Fanglomerates and landslide meg-
- abreccias crop out high in the section. North of Lake Havasu City a gently dipping latite flow (Tadf) dated by K-Ar at 19.9±0.5 Ma overlies more steeply tilted flows and strata (Tfts) from which an age of 19.2±0.5 Ma was obtained. The 30° and greater angular unconformity between the latite flow and underlying strata records the tilting of the Crossman block (Nielson and Beratan, 1988), so the age of tilting can be interpreted as near 19.5 Ma. Stratal dips indicate that tilting of other blocks ranged from before to after deposition of the 18.5-Ma Peach Springs Tuff.
- A middle Miocene fanglomerate unit (Tfg) represents the first of two distinct periods of postdetachment alluviation. Capping it are rhyolite and olivine basalt flows (Tr, Tob) that were dated at 12.2 to 8.6 Ma (Suneson and Lucchitta, 1983; Nakata and others, 1990). They and the underlying fanglomerate are relatively undeformed compared to older strata, but in the southeastern part of the map area they are faulted and are tilted gently southwest, the same
- The Bouse Formation of late Miocene and Pliocene age was deposited mostly in quiet water along the valley that was later to be followed by the Colorado River. A sandy subaerial facies of the Bouse was deposited in the northern part of the Mohave Mountains area; a similar facies was mapped south of the Whipple Mountains by Dickey and others (1980). Undeformed upper Miocene to Quaternary piedmont sedimentary deposits record a second period of alluviation. These deposits are fanglomerates composed of unsorted to poorly sorted, coarse, angular to subangular clastic debris derived from the exposed pre-Tertiary and Tertiary rocks. The fanglomerates interfinger with river-laid sand and gravel along the Colorado River west of the study area. Most of the fanglomerate deposits are at least as old as Pleistocene and probably formed a series of large, coalescing alluvial fans (unit QTs<sub>1</sub>). The depositional surface later was nearly completely dissected, with levels of erosion generally below the soil profile, on the west side of the mountains where drainages flow directly into the Colorado River. Dissection, planation, and reworking is largely responsible for a series of inset terraces and washes veneered by succesively younger parts of the piedmont unit. Soils are progressively less well developed on younger units (Wilshire and Reneau, 1992).

## Tertiary Structure

- Structural relations in the Crossman block indicate that it was tilted as a largely coherent block. The lower part of the Tertiary section dips steeply to overturned. The tilt persists at deeper structural levels in the block, as shown by the consistent steep orientations of rotated Proterozoic diabase sheets, subparallel to the lower Tertiary section. The Tertiary Mohave Mountains dike swarm maintains a moderate northeast dip all across the block and is unbroken by faults except in a few places.
- The evidence for tilting of the Crossman block implies that it exposes a cross section that represents progressively greater crustal depths from southwest to northeast. This section represents 10 to 15 km of crustal thickness if the Crossman block is unbroken and tilted at least 75° (Howard and John, 1987). Amphibolite bodies within the block trace cryptic Proterozoic folds that argue against much internal disruption of the block by Tertiary faulting, except where faults are mapped. The Crossman block may have been translated 40 km from Chemehuevi Valley, where the western Whipple Mountains below the Whipple Mountains detachment fault expose a footwall of similar dike swarms and Proterozoic gneisses (Carr and others, 1980; Dickey and others 1980; Davis and others, 1980, 1982; Howard and others, 1982a; Davis, 1988; Lister and Davis, 1989).
- The Whipple Mountains-Chemehuevi detachment fault projects eastward beneath the Crossman block and is inferred to juxtapose it against deeper mid-crustal mylonitic gneisses (see cross sections). In the eastern Chemehuevi Mountains and eastern Whipple Mountains the footwall of the detachment exposes mylonitic gneisses derived from Proterozoic to Tertiary protoliths and mulonitized in the Cretaceous and Tertiary (Anderson and Rowley, 1981: Davis and others, 1982; Anderson and others, 1988; Davis, 1988; John and Mukasa, 1990). An abrupt northeast termination of the Crossman block at its originally deep end is proposed on cross section A-A' in order to help explain a steep gravity gradient at this position. The gravity gradient led Simpson and others (1986) to model a steep boundary between denser rocks underlying the exposed Crossman block against less dense (lacking dikes?) bedrock to the northeast, under a thin sedimentary cover. The gravity model requires a greater density con-
- man block (Simpson and others, 1986). Hence the interpretation portrayed on cross section **A–A**' must be considered tentative and uncertain. A curving family of northwest-striking faults including the Wing fault cut the northwestern part of the Crossman block. These faults are steep, they cut the Mohave Mountains dike swarm, and they appear to have significant pre-dike-swarm displacements of gneiss units. Based on the generalized crustal section inferred for unbroken parts of the Crossman block, each fault of the Wing-fault family appears to attenuate crustal section, juxtaposing originally shallower rocks against originally deeper rocks to the north. Chloritized rock is present on the "deep" northeast side of the Wing fault but not on the "shallow" side, a pattern reminiscent of chloritized footwalls of low-angle normal faults in the region. Based on the resemblances to

trast (0.12 to 0.15 g/cm<sup>3</sup>) between bedrock terranes than would be expected from the largely

granitic bedrock exposed in the region and from density measurements of rocks in the Cross-

nally smaller dip. The Boulder Mine and Standard Wash areas expose numerous fault slices in synformal arrays structurally over the Crossman, Bill Williams, and Tumarion blocks. The faults in the Aubrey Hills and the Boulder Mine area mostly dip at low angles, and those in the Standard Wash area are inferred to also. Fault traces concave to the northeast in the Standard Wash area are consistent with synformal shapes and low northeast dips for the faults. Cross sections **B-B'** and **C-C'** interpret a shallow depth to the underlying footwall of the Whipple Mountains detachment fault. The depth is consistent not only with the geology but with the depth to a seismic-velocity increase interpreted as the base of the highly faulted allochthons (John and oth-

low-angle normal faults in the region, the steep Wing fault perhaps was rotated from an origi-

- ers, 1987; McCarthy and others, 1987). The southeast-dipping Powell Peak fault bounds the Tumarion block against higher blocks of the Boulder Mine area, and the southeast-dipping Crossman Peak fault similarly bounds the exposed Crossman block against higher blocks of the Standard Wash area. Measured striae on the latter fault strike northeast in a direction parallel to the direction of extension and regional fault slips, including slip on the deeper Whipple Mountains and Chemehuevi detachment faults (Davis and others, 1980; Howard and John, 1987; John, 1987a).
- Cross sections **B-B**' and **C-C**' portray the Powell Peak and Crossman Peak faults as bottoming at shallow depth, so that the Crossman block is connected at depth below the Standard Wash area and the Boulder Mine area with the Bill Williams and Tumarion blocks. This interpretation is based on (1) the alignment of Tertiary strata in the three big upended blocks; (2) the moderate (30-40°) dip of the exposed Powell Peak and Crossman Peak faults; and (3) the synformal array of higher blocks in the Boulder Mine and Standard Wash areas (Howard and others, 1982a; Howard and John, 1987). An alternate interpretation is that the Crossman Peak and Powell Peak faults are syndepositional transfer faults, and that the Crossman block does not intervene between the Standard Wash area and the Aubrey Hills as portrayed on the cross sections. This latter interpretation is based on stratigraphic coherence in the Tertiary section of the Aubrey Hills and Standard Wash areas and stratigraphic contrasts to the Crossman block (Beratan and others, 1990; Nielson and Beratan, 1990).
- Relatively mild tilting and step faulting in upper Miocene strata (northeastern part of cross section B-B') may postdate active extension accommodated on the Whipple Mountains and Crossman Peak faults. The late Miocene tilting may relate to rebound of the detachment faults and later adjustments of tilted blocks following the end of movement on the Whipple Mountains

# ACKNOWLEDGMENTS

Sherrod, R.W. Tabor, and J.G. Evans led to many improvements.

# REFERENCES CITED

- ris, L.G., Jr., Byers, C.W., Mickelson, D.M., and Shanks, W.C., eds., Proterozoic Geology; Selected Papers from an International Proterozoic Symposium: Geological Society of Anderson, J.L., and Bender, E.E., 1989, Nature and origin of Proterozoic A-type granitic magmatism in the southwestern United States of America: Lithos, v. 23, p. 19–52.
- ated metaluminous granitoids, Whipple Mountains, California: Canadian Mineralogist, v. 19, p. 83–101.
- Southern California Edison Co., 66 p.
- Burchfiel, B.C., and Davis, G.A., 1981, Mojave desert and environs, in Ernst, W.G., ed., The Carr, W.J., 1991, A contribution to the structural history of the Vidal-Parker region, California

- MISCELLANEOUS INVESTIGATIONS SERIES MAP I-2308 (SHEET 2 OF 2) Davis, G.A., 1988, Rapid upward transport of mid-crustal mylonitic gneisses in the footwall of
- a Miocene detachment fault, Whipple Mountains, southeastern California: Geologische Rundschau, v. 77, p. 191–209. Davis, G.A., Anderson, J.L., Frost, E.G., and Shackelford, T.J., 1980, Mulonitization and detachment faulting in the Whipple-Buckskin-Rawhide Mountains terrane, southeastern California and western Arizona, in Crittenden, M.D., Jr., and others, eds., Cordilleran Metamorphic Core Complexes: Geological Society of America Memoir 153, p. 79–129.
- Davis, G.A., Anderson, J.L., Martin, D.L., Krummenacher, Daniel, Frost, E.G., and Arm strong, R.L., 1982, Geologic and geochronologic relations in the lower plate of the Whipple detachment fault, Whipple Mountains, southeastern California: A progress report, in Frost, E.G., and Martin, D.L., eds., Mesozoic-Cenozoic Tectonic Evolution of the Colorado River Region, California, Arizona, and Nevada: San Diego, Calif., Cordilleran Publishers, p. 408–432.
- Davis, G.A., and Lister, G.S., 1988, Detachment faulting in continental extension: Perspectives from the southwestern U.S. Cordillera, in Clark, S.P., Jr., Burchfiel, B.C., and Suppe John, eds., Processes in continental lithospheric deformation: John Rodgers symposium volume: Geological Society of America Special Paper 218, p. 133–159. Dickey, D.D., Carr, W.J., and Bull, W.B., 1980, Geologic map of the Parker NW, Parker, and
- parts of the Whipple Mountains SW and Whipple Wash quadrangles, California and Arizona, U.S. Geological Survey Miscellaneous Investigations Series Map I-1124, scale Dunn, J.F., 1986, The structural geology of northeastern Whipple Mountains detachment terrane: Los Angeles, University of Southern California, M.S. thesis, 113 p. Fitzgibbon, T.T., 1988, Tectonic significance and characteristics of Middle Proterozoic diabase
- sheets, eastern Mojave Desert region, California and Arizona [abs.]: Geological Society of America Abstracts with Programs, v. 20, no. 3, p. 160. Fitzgibbon, T.T., and Howard, K.A., 1987, Tectonic significance of Middle Proterozoic diabase sheets in southeastern California and Arizona [abs.]: Geological Society of America
- Abstracts with Programs, v. 19, no. 6, p. 377. Foster, D.A., Harrison, T.M., Miller, C.F., and Howard, K.A., 1990, The  $^{40}\text{Ar}/^{39}\text{Ar}$  thermochronology of the eastern Mojave Desert, California, and adjacent western Arizona with implications for the evolution of metamorphic core complexes: Journal of Geophysical Research, v. 95, p. 20,005–20,024.
- Glazner, A.F., Nielson, J.E., Howard, K.A., and Miller, D.M., 1986, Correlation of the Peach Springs Tuff, a large-volume Miocene ignimbrite sheet in California and Arizona: Geology, v. 14. p. 840–843. Hendricks, J.D., and Lucchitta, I., 1974, Upper Precambrian igneous rocks of the Grand Canvon, Arizona, in Karlstrom, N.V., and Swann, G.A., eds., Geology of Northern Arizona, Part I, Regional studies: Geological Society of America, Rocky Mountain Section Meeting,
- Flagstaff, Arizona, p. 65–86. Howard, K.A., 1991, Intrusion of horizontal dikes: Tectonic significance of Middle Proterozoic
- diabase sheets widespread in the upper crust of the southwestern United States: Journal of Geophysical Research, v. 96, p. 12,461–12,478. Howard, K.A., Goodge, J.W., and John, B.E., 1982a, Detached crystalline rocks of the Mohave, Buck, and Bill Williams Mountains, western Arizona, in Frost, E.G., and Martin, D.L., eds., Mesozoic-Cenozoic Tectonic Evolution of the Colorado River Region, California, Arizona, and Nevada: San Diego, Calif., Cordilleran Publishers, p. 377–390. Howard, K.A., and John, B.E., 1987, Crustal extension along a rooted system of low-angle
- normal faults, Colorado River extensional corridor, California and Arizona, in Coward, M.D., Dewey, S.F., and Hancock, P.L., eds., Continental extensional tectonics: Geological Society of London Special Publication 28, p. 299–311. Howard, K.A., John, B.E., and Miller, C.F., 1987, Metamorphic core complexes, Mesozoic ductile thrusts, and Cenozoic detachments: Old Woman Mountains-Chemehuevi Mountains transect, California and Arizona, in Davis G.H., and VanderVolder, E.M., eds., Geologic diversity of Arizona and its margins: Excursion to choice areas: Arizona Bureau of
- Geology and Mineral Technology, Geological Survey Branch, Special Paper 5, p. Howard, K.A., Stone, Paul, Pernokas, M.A., and Marvin, R.F., 1982b, Geologic and geochronologic reconnaissance of the Turtle Mountains area, California; west border of the Whipple detachment terrane, in Frost, E.G., and Martin, D.L., eds., Mesozoic-Cenozoic Tectonic Evolution of the Colorado River Region, California, Arizona, and Nevada: San
- Diego, Calif., Cordilleran Publishers, p. 341–354. John, B.E., 1981, Reconnaissance study of Mesozoic plutonic rocks in the Mojave Desert region. in Howard, K.A., Carr, M.D., and Miller, D.M., eds., Tectonic framework of the Mojave and Sonoran Deserts, California and Arizona: U.S. Geological Survey Open-File Report 81–503, p. 48–50.
- John, B.E., 1987a, Geometry and evolution of a mid-crustal extensional fault system: Chemehuevi Mountains, southeastern California, in Coward, M.D., Dewey, J.F., and Hancock P.L., eds., Continental Extensional Tectonics: Geological Society of London Special Publi cation 28, p. 313–335.
- John, B.E., 1987b, Geologic map of the Chemehuevi Mountains area, San Bernardino County, California and Mohave County, Arizona: U.S. Geological Survey Open-File Report 87–666, scale 1:24,000. John, B.E., 1988, Structural reconstruction and zonation of a tilted mid-crustal magma cham
- ber: The felsic Chemehuevi Mountains plutonic suite: Geology, v. 16, p. 613–617. John, B.E., Howard, K.A., and McCarthy, J., 1987, Geologic and geophysical evidence for the nature of upper and mid-crustal continental extension in the Colorado River area, California and Arizona [abs.]: Geological Society of America Abstracts with Programs, v. 19,
- John, B.E., and Mukasa, S.B., 1990, Footwall rocks to the mid-Tertiary Chemehuevi detachment fault: A window into the middle crust in the southern Cordillera: Journal of Geophysical Research, v. 95, p. 463–486. John, B.E., and Wooden, J., 1990, Petrology and geochemistry of the metaluminous to pera-
- luminous Chemehuevi Mountains Plutonic Suite, southeastern California, in Anderson, J.L., ed., The nature and origin of Cordilleran magmatism: Geological Society of America Memoir 174, p. 71–98. Light, T.D., and McDonnell, J.R., Jr., 1983, Mineral investigation of the Crossman Peak Wilderness Study Area, Mohave County, Arizona: U.S. Bureau of Mines MLA 82–83, 203 p. Light, T.D., Pike, J.E., Howard, K.A., McDonnell, J.R., Jr., Simpson, R.W., Raines, G.L.
- Knox, R.D., Wilshire, H.G., and Pernokas, M.A., 1983, Mineral resource potential of the Crossman Peak Wilderness Study Area (5–7B), Mohave County, Arizona: U.S. Geological Survey Miscellaneous Field Studies Map MF-1602-A, 21 p., scale 1:48,000. Lister, G.S., and Davis, G.A., 1989, The origin of metamorphic core complexes and detach-
- ment faults formed during Tertiary continental extension in the northern Colorado River region, U.S.A.: Journal of Structural Geology, v. 11, p. 65–94. Lucchitta, Ivo, and Suneson, Neil, 1988, Geologic map of the Planet 2 SW quadrangle Mohave County, Arizona: U.S. Geological Survey Open-file Report 88-547, scale
- Lucchitta, Ivo, and Suneson, N.H., 1994, Geologic map of the Castaneda Hills quadrangle Mohave County, Arizona: U.S. Geological Survey, Geological Survey Map GQ-1720, Lucchitta, Ivo and Suneson, N.H., 1994, Geologic map of the Castaneda Hills SW quadrangle
- Mohave County, Arizona: U.S. Geological Survey, Geologic Quadrangle Map GQ-1719, Lucchitta, Ivo, and Suneson, N.H., 1994, Geologic map of the Centennial Wash quadrangle Mohave County, Arizona: U.S. Geological Survey, Geologic Quadrangle, Map GQ-1718,
- McCarthy, J., Fuis, G.S., and Wilson, J., 1987, Crustal structure in the region of the Whipple Mountains metamorphic core complex, from seismic refraction results [abs.]: Geological Society of America Abstracts with Programs, v. 19, n. 7, p. 763. Nakata, J.K., 1982, Preliminary report on diking events in the Mohave Mountains, Arizona in
- Frost, E.G. and Martin, D.L., eds., Mesozoic-Cenozoic Tectonic Evolution of the Colorado River Region, California, Arizona, and Nevada: San Diego, Calif., Cordilleran Publishers, Nakata, J.K., Pernokas, M.A., Howard, K.A., Nielson, J.E., and Shannon, J.R., 1990, K-Ar
- and fission-track ages (dates) of volcanic, intrusive, altered, and metamorphic rocks in the Mohave Mountains area, west-central Arizona: Isochron/West, no. 56, p. 8–20. Nielson, J.E., 1986, Miocene stratigraphy of the Mohave Mountains, Arizona, and correlation with adjacent ranges, in Cenozoic Stratigraphy, Structure, and Mineralization in the Mojave Desert: Geological Society of America Cordilleran Section, 82nd Annual Meeting
- Guidebook and volume, Trips 5 and 6, p. 15–24. Nielson, J.E., and Beratan, K.K., 1988, Time of detachment faulting from Miocene sequences Colorado River extended terrane [abs.]: Geological Society of America Abstracts with Programs, v. 20, p. 393. Nielson, J.E., and Beratan, K.K., 1990, Tertiary basin development and tectonic implications,
- Whipple detachment system, Colorado River extensional corridor, California and Arizona: Journal of Geophysical Research, v. 95, p. 599–614. Nielson, J.E., Lux, D.R., Dalrymple, G.B., and Glazner, A.F., 1990, Age of the Peach Springs Tuff: Journal of Geophysical Research, v. 95, p. 571–580. Pike, J.E.N., and Hansen, V.L., 1982, Complex Tertiary stratigraphy and structure, Mohave
- Mountains, Arizona: A preliminary report, in Frost, E.G., and Martin, D.L., eds, Mesozoic-Cenozoic Tectonic Evolution of the Colorado River Region, California, Arizona, and Nevada: San Diego, Calif., Cordilleran Publishers, p. 91–96. Rehrig, W.A., and Heidrick, T.L., 1976, Regional tectonic stress during the Laramide and late Tertiary intrusive periods, Basin and Range province, Arizona: Arizona Geological Society
- Digest. v. 10. p. 205–228. Sherrod, D.R., 1988, Preliminary geologic map of the Monkey's Head 7.5-minute quadrangle Mohave County, Arizona: U.S. Geological Survey Open-File Report 88-899, scale
- Simpson, R.W., Gage, T.B., and Bracken, R.E., 1986, Aeromagnetic and isostatic gravity maps of the Crossman Peak Wilderness Study Area, Mohave County, Arizona: U.S. Geological Survey Miscellaneous Field Studies Map MF-1602-B, scale 1:48,000. Streckeisen, A.L., 1973, Plutonic rocks: Classification and nomenclature recommended by the IUGS Subcommission on the Systematics of Igneous Rocks: Geotimes, v. 18, no. 10, p.
- Suneson, N.H., 1980, The origin of bimodal volcanism, west-central Arizona: Santa Barbara, University of California, Ph.D. dissertation, 293 p. Suneson, N.H., and Lucchitta, Ivo, 1983, Origin of bimodal volcanism, southern Basin and Range province, west-central Arizona: Geological Society of America Bulletin, v. 94, p.
- Wilshire, H.G., and Reneau, S.L., 1992, Geomorphic surfaces and underlying deposits of the Mohave Mountains piedmont, lower Colorado River, Arizona: Zeitschrift für Geomorphologie, v. 36, p. 207–226. Wilson, E.D., and Moore, R.T., 1959, Geologic map of Mohave County, Arizona: Arizona Bureau of Mines, University of Arizona, Tucson, scale 1:375,000. Wrucke, C.T., and Shride, A.R., 1972, Correlation of Precambrian diabase in Arizona and
- Young, R.A., and Brennan, W.J., 1974, Peach Springs Tuff: Its bearing on structural evolution of the Colorado Plateau and development of Cenozoic drainage in Mohave County, Arizona: Geological Society of America Bulletin, v. 85, p. 83–90. Wooden, J.L., and Miller, D.M., 1990, Chronologic and isotopic framework for Early Proterozoic crustal evolution in the eastern Mojave Desert region, SE California: Journal of Geo-

physical Research, v. 95, p. 20,133–20,146.

southern California [abs.]: Geological Society America Abstracts with Programs, v. 4, p.

GEOLOGIC MAP OF THE MOHAVE MOUNTAINS AREA, MOHAVE COUNTY, WESTERN ARIZONA