

Math 3298 Practice final exam problems

The actual exam will consist of six to eight required questions and possibly an optional extra credit question.

- (1) Reverse the order of integration for the integral  $\int_0^1 \int_x^1 \int_0^{y^2} f(x, y, z) dz dy dx$ .
- (2) Compute the value of the vector line integral  $\int_C \vec{F} \cdot d\vec{r}$  where  $C$  is the path  $(4 - 3^t, -2 + 2t, \pi t)$ ,  $t \in [0, 1]$ , and  $\vec{F} = (2x \cos z - x^2, z - 2y, y - x^2 \sin z)$ .
- (3) Find the linearization of  $f(x, y)$  at  $(x, y) = (0, 1)$  if  $f = h(u(x, y), v(x, y))$  and  $\text{grad}(h)|_{(1,1)} = (\frac{\partial h}{\partial u}, \frac{\partial h}{\partial v})|_{(1,1)} = (2, 3)$ ,  $u(x, y) = x + y$ , and  $v(x, y) = y^2$ .
- (4) Find the surface area of the torus parameterized by  $x = (2 + \cos(v)) \cos(u)$ ,  $y = (2 + \cos(v)) \sin(u)$ ,  $z = \sin(v)$ , with  $u \in [0, 2\pi]$  and  $v \in [0, 2\pi]$ .
- (5) Find the maxima and minima of  $f(x, y) = \frac{1}{x} + \frac{2}{y}$  on the set  $\frac{1}{x^2} + \frac{1}{y^2} = 1$ .
- (6) Find the volume of the solid wedge bounded by the planes  $z = 0$  and  $z = -2y$  and the cylinder  $x^2 + y^2 = 4$  (with  $y \geq 0$ ).
- (7) Use Green's Theorem to find the smooth, simple, closed and positively oriented curve in the plane for which the line integral  $\oint (\frac{x^2 y}{4} + \frac{y^3}{3}) dx + x dy$  has the largest possible value.
- (8) Compute the value of  $\int \int_S (\nabla \times \vec{F}) \cdot \vec{n} dS$  where  $S$  is the upper half of the ellipsoid  $4x^2 + 9y^2 + 36z^2 = 36$ ,  $z \geq 0$ , with upward pointing normal, and  $\vec{F} = (y, x^2, (x^2 + y^2)^{3/2} e^{xyz})$ .
- (9) Let  $\vec{r}(t)$  be a curve in space with unit tangent, normal, and binormal vectors  $\vec{T}$ ,  $\vec{N}$ , and  $\vec{B} = \vec{T} \times \vec{N}$ . Show that  $\frac{d\vec{B}}{dt}$  is perpendicular to  $\vec{T}$ .
- (10) Compute the flux integral  $\int \int_S \vec{F} \cdot \vec{n} dS$  where  $S$  is the graph of  $z = 1 - x^2 - y^2$ , with upward normal, for  $z \geq 0$ , and with  $\vec{F} = (xz, yz, 2z^2)$ .
- (11) Use the divergence theorem to compute the flux of  $\vec{F} = (z^5 + x, \cos(xz), z^2)$  through the surface bounded by  $z = 0$  and  $z = 1 - x^2 - y^2$ .